



TOHOKU  
UNIVERSITY

# Magnetic Fields Evolution in Grand-design Spiral Galaxies with 3D MHD Simulations

Sho Nakamura (Tohoku Univ.)






TOHOKU  
UNIVERSITY


# Introduction

# Spiral Arms

# Spiral Arms





# Grand-design Spiral Galaxies



Fletcher+, 2011  
(M51,  $\lambda=6\text{cm}$ )

Spiral galaxies have magnetic fields along the spiral arms  
@ galactic plane.

# Grand-design Spiral Galaxies



Fletcher+, 2011

Structure of the magnetic fields are different in the disc and halo.



TOHOKU  
UNIVERSITY

# Motivation

# Motivation

Effects of a spiral potential on

1. gaseous structure
- 2. magnetic field structure**

# Model

&

# Methods



TOHOKU  
UNIVERSITY



# Model: Basic Equations

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = 0$$

$$\frac{\partial}{\partial t}(\rho \mathbf{u}) + \nabla \cdot \left\{ \rho \mathbf{u} \mathbf{u} + \left( p_{\text{gas}} + \frac{1}{8\pi} B^2 \right) \mathbf{I} - \frac{1}{4\pi} \mathbf{B} \mathbf{B} \right\} = -\rho \nabla \Phi + \rho \Omega_{\text{sp}}^2 \mathbf{R} - 2\rho \boldsymbol{\Omega}_{\text{sp}} \times \mathbf{u}$$

$$\frac{\partial e}{\partial t} + \nabla \cdot \left[ \left\{ e + \left( p_{\text{gas}} + \frac{1}{8\pi} B^2 \right) \right\} \mathbf{u} - \frac{1}{4\pi} (\mathbf{B} \cdot \mathbf{u}) \mathbf{B} \right] = \mathbf{u} \cdot (-\rho \nabla \Phi + \rho \Omega_{\text{sp}}^2 \mathbf{R}) - \rho^2 \Lambda(T_{\text{gas}})$$

$$\frac{\partial \mathbf{B}}{\partial t} - \nabla \cdot (\mathbf{u} \mathbf{B} - \mathbf{B} \mathbf{u}) = \mathbf{0}$$

$$e_{\text{gas}} = \frac{p_{\text{gas}}}{\gamma_{\text{gas}} - 1}, \quad \gamma_{\text{gas}} = 5/3, \quad e = \frac{1}{2} \rho u^2 + e_{\text{gas}} + \frac{1}{8\pi} B^2, \quad \mathbf{u} = \mathbf{v} - \Omega_{\text{sp}} R \mathbf{e}_\varphi$$

# Model: Basic Equations

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = 0$$


$$\frac{\partial}{\partial t}(\rho \mathbf{u}) + \nabla \cdot \left\{ \rho \mathbf{u} \mathbf{u} + \left( p_{\text{gas}} + \frac{1}{8\pi} B^2 \right) \mathbf{I} - \frac{1}{4\pi} \mathbf{B} \mathbf{B} \right\} = -\rho \nabla \Phi + \rho \Omega_{\text{sp}}^2 \mathbf{R} - 2\rho \boldsymbol{\Omega}_{\text{sp}} \times \mathbf{u}$$

$$\frac{\partial e}{\partial t} + \nabla \cdot \left[ \left\{ e + \left( p_{\text{gas}} + \frac{1}{8\pi} B^2 \right) \right\} \mathbf{u} - \frac{1}{4\pi} (\mathbf{B} \cdot \mathbf{u}) \mathbf{B} \right] = \mathbf{u} \cdot (-\rho \nabla \Phi + \rho \Omega_{\text{sp}}^2 \mathbf{R}) - \rho^2 \Lambda(T_{\text{gas}})$$

$$\frac{\partial \mathbf{B}}{\partial t} - \nabla \cdot (\mathbf{u} \mathbf{B} - \mathbf{B} \mathbf{u}) = \mathbf{0}$$

$$e_{\text{gas}} = \frac{p_{\text{gas}}}{\gamma_{\text{gas}} - 1}, \quad \gamma_{\text{gas}} = 5/3, \quad e = \frac{1}{2} \rho u^2 + e_{\text{gas}} + \frac{1}{8\pi} B^2, \quad \mathbf{u} = \mathbf{v} - \Omega_{\text{sp}} R \mathbf{e}_\varphi$$

# Radiative Loss Function



Raymond+, 1979

$$\Lambda(T_{\text{gas}}) = \begin{cases} 0 & (T_{\text{gas}} < 2 \times 10^4) \\ 10^{-21.85} & (2 \times 10^4 < T_{\text{gas}} < 4 \times 10^4) \\ 10^{-31} T_{\text{gas}}^2 & (4 \times 10^4 < T_{\text{gas}} < 7.9 \times 10^4) \\ 10^{-21.2} & (7.9 \times 10^4 < T_{\text{gas}} < 2.5 \times 10^5) \\ 10^{-10.4} T_{\text{gas}}^{-2} & (2.5 \times 10^5 < T_{\text{gas}} < 5.6 \times 10^5) \\ 0 & (5.6 \times 10^5 < T_{\text{gas}}) \end{cases}$$

Rosner+, 1978

# Model: Basic Equations

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = 0$$

$$\frac{\partial}{\partial t}(\rho \mathbf{u}) + \nabla \cdot \left\{ \rho \mathbf{u} \mathbf{u} + \left( p_{\text{gas}} + \frac{1}{8\pi} B^2 \right) \mathbf{I} - \frac{1}{4\pi} \mathbf{B} \mathbf{B} \right\} = -\rho \nabla \Phi + \rho \Omega_{\text{sp}}^2 \mathbf{R} - 2\rho \boldsymbol{\Omega}_{\text{sp}} \times \mathbf{u}$$

$$\frac{\partial e}{\partial t} + \nabla \cdot \left[ \left\{ e + \left( p_{\text{gas}} + \frac{1}{8\pi} B^2 \right) \right\} \mathbf{u} - \frac{1}{4\pi} (\mathbf{B} \cdot \mathbf{u}) \mathbf{B} \right] = \mathbf{u} \cdot (-\rho \nabla \Phi + \rho \Omega_{\text{sp}}^2 \mathbf{R}) - \rho^2 \Lambda(T_{\text{gas}})$$

$$\frac{\partial \mathbf{B}}{\partial t} - \nabla \cdot (\mathbf{u} \mathbf{B} - \mathbf{B} \mathbf{u}) = \mathbf{0}$$

$$e_{\text{gas}} = \frac{p_{\text{gas}}}{\gamma_{\text{gas}} - 1}, \quad \gamma_{\text{gas}} = 5/3, \quad e = \frac{1}{2} \rho u^2 + e_{\text{gas}} + \frac{1}{8\pi} B^2, \quad \mathbf{u} = \mathbf{v} - \Omega_{\text{sp}} R \mathbf{e}_\varphi$$

# Spiral Potential

$$\Phi(R, \varphi, z; t) = \Phi_{\text{bulge}}(R, z) + \Phi_{\text{disc}}(R, z) + \Phi_{\text{halo}}(R, z) + \Phi_{\text{sp}}(R, \varphi, z; t)$$

**non-axisymmetric potential**

$$\Phi_{\text{sp}}(R, \varphi, z; t) = \Phi_{\text{disc}}(R, z) \epsilon_{\text{sp}} \frac{(R/R_a)^2}{\{1 + (R/R_a)^2\}^{3/2}} \frac{z_a}{\sqrt{z^2 + z_a^2}} \times \cos \left[ m \left\{ \varphi - \Omega_{\text{sp}} t + \cot i_{\text{sp}} \ln \left( \frac{R}{R_{\text{phase}}} \right) \right\} \right]$$

Wada+ 2004, Baba+ 2011

$$\epsilon_{\text{sp}} = 0.025, R_a = 7\text{kpc}, z_a = 0.3\text{kpc}, m = 2, \Omega_{\text{sp}} = 15\text{kms}^{-1}\text{kpc}^{-1}, i_{\text{sp}} = 15^\circ, R_{\text{phase}} = 0.1\text{kpc}$$


# Model: Initial Condition

Initial gas distribution

$$\rho = \rho_{\text{disc}} + \rho_{\text{halo}}, \quad \rho_{\text{disc}} = \rho_{\text{disc},0} \exp\left(-\frac{R}{R_{\text{disc}}}\right) \text{sech}^2\left(\frac{z}{z_{\text{disc}}}\right)$$
$$\rho_{\text{halo}} = \rho_{\text{halo},0} \exp\left(-\frac{\Phi_{\text{axi}}(R, z) - \Phi_{\text{axi}}(0, 0)}{C s_{\text{halo}}^2}\right)$$

$$R_{\text{disc}} = 4\text{kpc}, z_{\text{disc}} = 0.3\text{kpc}, T_{\text{disc}} = 10^4\text{K}, T_{\text{halo}} = 10^6\text{K}$$

Rotation curve



# Model: Initial Condition

Initial magnetic field distribution

$$B_R = B_z = 0, B_\varphi = B_{\varphi,0} \exp\left(-\frac{R}{R_{\text{disc}}}\right) \text{sech}^2\left(\frac{z}{z_{\text{disc}}}\right)$$

$$\beta = 10^4 \text{ @ } R = 0\text{kpc} \implies B_{\varphi,0} \sim 0.1\mu\text{G}$$

# Methods

I. flux: HLLD (Miyoshi&Kusano 2005)

II. time & space: 2nd order accuracy

III. divergence constraint:

Hyperbolic Divergence Cleaning (Dedner+ 2002)

IV. cylindrical coordinate,  $(N_R, N_\varphi, N_z)=(250, 64, 200)$

V.  $0.2\text{kpc} < R < 21.9\text{kpc}$ ,  $0 < \varphi < 2\pi$ ,  $0 < z < 2.4\text{kpc}$






TOHOKU  
UNIVERSITY


# Results

# Results

Log(density)





Log(temperature)



# Results( $t=800\text{Myr}$ )

Log(density)



Log(temperature)



# Results

$\text{Log}(B^2/8\pi)$



$B_\varphi(\mu\text{G})$



# Results( $t=800\text{Myr}$ )


$\text{Log}(B^2/8\pi)$

$B_\varphi(\mu\text{G})$




# Results

$B_y(\mu\text{G})$



# Results( $t=800\text{Myr}$ )



$B_y(\mu\text{G})$



# Results( $t=800\text{Myr}$ )

$\text{Log}(B^2/8\pi)$


$B_\phi(\mu\text{G})$



?






# MRI-Parker dynamo




Nishikori+, 2006  
Machida+, 2013

# Results( $t=800\text{Myr}$ )



# Results

time evolution of plasma  $\beta$  ( $4\text{kpc} < R < 6\text{kpc}$ ,  $0 < z < 0.15\text{kpc}$ )





TOHOKU  
UNIVERSITY

# Conclusion

# Conclusion

1. Due to the “MRI-Parker dynamo” process, toroidal component of magnetic field change reversal in disc region.
2. We have shown that spiral arms rapidly amplify magnetic fields.



TOHOKU  
UNIVERSITY

Thank you  
for your attention

**Dziękuję**